Recombinant Human CSNK2A1/CK2A1 Protein (GST Tag)

Catalog Number: PKSH030405



Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Synonyms CK2A1;CKII;CSNK2A3

Species Human

Expression Host Baculovirus-Insect Cells

SequenceMet 1-Gln 391AccessionNP_808227.1Calculated Molecular Weight71.4 kDaObserved molecular weight65 kDaTagN-GST

Bioactivity The specific activity was determined to be 9 nmol/min/mg using casein as substrate.

Properties

Purity > 93 % as determined by reducing SDS-PAGE.

Endotoxin < 1.0 EU per µg of the protein as determined by the LAL method.

Storage Storage Store at $< -20^{\circ}$ C, stable for 6 months. Please minimize freeze-thaw cycles.

Shipping This product is provided as liquid. It is shipped at frozen temperature with blue


ice/gel packs. Upon receipt, store it immediately at < - 20°C.

Formulation Supplied as sterile solution of 50mM Tris, 100mM NaCl, 0.5mM PMSF, 0.5mM

GSH, pH 8.0

Reconstitution Not Applicable

Data



> 93 % as determined by reducing SDS-PAGE.

Background

Casein kinase II subunit alpha, also known as CK II alpha, CSNK2A1 and CK2A1, is a member of the protein kinase superfamily, Ser / Thr protein kinase family and CK2 subfamily. Casein kinase II (CSNK2A1) is a serine / threonine protein kinase that phosphorylates acidic proteins such as casein. This kinase is composed of an alpha, an alpha-prime, and two beta subunits. The alpha subunits contain the catalytic activity while the beta subunits undergo autophosphorylation. Casein kinase II (CSNK2A1) is a constitutively active, ubiquitously expressed serine / threonine protein kinase that is thought to have a regulatory function in cell proliferation, cell differentiation and apoptosis. CSNK2A1 functions as a tetrameric complex consisting of two regulatory beta-subunits and two catalytic units (alpha and alpha') in a homomeric or heteromeric conformation. Whilst the alpha- and alpha'-subunits are catalytically identical, proteins that regulate CSNK2A1, such as cdc2 and Hsp90, preferentially bind to the alpha and not the alpha'-subunit. CSNK2A1 can phosphorylate a number of key intracellular signaling proteins implicated in tumor suppression (p53 and

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: <u>www.elabscience.com</u> Email: <u>techsupport@elabscience.com</u>

Recombinant Human CSNK2A1/CK2A1 Protein (GST Tag)

Catalog Number:PKSH030405



PTEN) and tumorigenesis (myc, jun, NF-kappaB). CSNK2A1 is also thought to influence Wnt signaling via beta-catenin phosphorylation and the PI 3-K signaling pathway via th phosphorylation of Akt.

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: <u>www.elabscience.com</u> Email: <u>techsupport@elabscience.com</u>