

## Recombinant Human PDGF-BB Protein (His Tag)

Catalog No. PKSH030442

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Synonyms</b>	Platelet-Derived Growth Factor Subunit B;PDGF Subunit B;PDGF-2;Platelet-Derived Growth Factor B Chain;Platelet-Derived Growth Factor Beta Polypeptide;Proto-Oncogene c-Sis;Becaplermin;PDGFB;PDGF2;SIS
<b>Species</b>	Human
<b>Expression Host</b>	Yeast
<b>Sequence</b>	Ser82-Thr190
<b>Accession</b>	NP_148937.1
<b>Calculated Molecular Weight</b>	14.3 kDa
<b>Tag</b>	N-His
<b>Bioactivity</b>	Immobilized human PDGFB at 10µg/mL (100µL/well) can bind human PDGFRβ-Fch, the EC50 of human PDGFRβ-Fch is 0.05-0.2 µg/mL.

### Properties

<b>Purity</b>	> 95 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	Please contact us for more information.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from sterile PBS, 25 % Glycerol, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01 % Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

### Data



> 95 % as determined by reducing SDS-PAGE.

### Background

#### For Research Use Only

Platelet-Derived Growth Factor Subunit B (PDGFB) belongs to the PDGF/VEGF growth factor family. Platelet-derived growth factor is a potent mitogen for cells of mesenchymal origin. PDGFB can exist either as a homodimer (PDGF-BB) or as a heterodimer with the platelet-derived growth factor alpha polypeptide (PDGF-AB), where the dimers are connected by disulfide bonds. Mutations in this gene are associated with meningioma. Binding of PDGFB to its receptor elicits a variety of cellular responses. In addition, PDGFB is released by platelets upon wounding and plays an important role in stimulating adjacent cells to grow and thereby heals the wound.