

Recombinant Human RHEB Protein (His Tag)

Catalog Number:PKSH031137



Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

| | |
|------------------------------------|---|
| Synonyms | GTP-Binding Protein Rheb;Ras Homolog Enriched in Brain;RHEB;RHEB2 |
| Species | Human |
| Expression Host | Baculovirus-Insect Cells |
| Sequence | Met 1-Cys181 |
| Accession | Q15382 |
| Calculated Molecular Weight | 22.4 kDa |
| Observed molecular weight | 21 kDa |
| Tag | N-His |

Properties

| | |
|-----------------------|--|
| Purity | > 85 % as determined by reducing SDS-PAGE. |
| Endotoxin | < 1.0 EU per µg of the protein as determined by the LAL method. |
| Storage | Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months. |
| Shipping | This product is provided as lyophilized powder which is shipped with ice packs. |
| Formulation | Lyophilized from sterile 20mM Tris, 500mM NaCl, 10% glycerol, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual. |
| Reconstitution | Please refer to the printed manual for detailed information. |

Data



> 85 % as determined by reducing SDS-PAGE.

Background

RHEB is a recently discovered member of the Ras superfamily that may be involved in neural plasticity. This function is novel and not typically associated with the Ras proteins. RHEB gene is a member of the small GTPase superfamily and encodes a lipid-anchored; cell membrane protein with five repeats of the RAS-related GTP-binding region. RHEB is vital in regulation of growth and cell cycle progression due to its role in the insulin / TOR / S6K signaling pathway. The protein has GTPase activity and shuttles between a GDP-bound form and a GTP-bound form; and farnesylation of RHEB is required for this activity. Three pseudogenes have been mapped; two on chromosome 10 and one on chromosome 22.

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: www.elabscience.com

Tel: 1-832-243-6086

Email: techsupport@elabscience.com

Fax: 1-832-243-6017