

Recombinant Mouse Wnt Inhibitory Factor 1/WIF1 Protein (His Tag)

Catalog No. PKSM040406

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	AW107799;WIF-1
Species	Mouse
Expression Host	HEK293 Cells
Sequence	Met1-Trp379
Accession	Q9WUA1
Calculated Molecular Weight	39.8 kDa
Observed molecular weight	44 kDa
Tag	C-His
Bioactivity	Not validated for activity

Properties

Purity	> 93 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 93 % as determined by reducing SDS-PAGE.

Background

WIF1, also known as WIF-1 and wnt inhibitory factor 1, is a secreted protein which binds WNT proteins and inhibits their activities. It contains a WNT inhibitory factor (WIF) domain and 5 epidermal growth factor (EGF)-like domains.

For Research Use Only

WNT proteins are extracellular signaling molecules involved in the control of embryonic development. WIF1 may be involved in mesoderm segmentation and can be detected in fish, amphibia and mammals. WIF-1 is a recurrent target in human salivary gland oncogenesis. Downregulation of WIF1 takes part in the development and progression of pleomorphic adenomas. WIF1 also is a tumor suppressor, and has been found to be epigenetically silenced in various cancers, specifically in nonfunctioning pituitary tumors. WIF1 are expected to have molecular function (protein tyrosine kinase activity) and to localize in various compartments (extracellular space, extracellular region).