

Recombinant Mouse FGFR5/FGFR5 Protein (His Tag)

Catalog Number:PKSM040824



Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	FGFR5;FGFR5beta;FGFR5gamma
Species	Mouse
Expression Host	HEK293 Cells
Sequence	Met 1-Pro 374
Accession	NP_473412.1
Calculated Molecular Weight	40.4 kDa
Observed molecular weight	55-60 kDa
Tag	C-His
Bioactivity	1. Immobilized mouse at 10 µg/ml (100 µl/well) can bind mouse FGFR5. The EC50 of mouse FGFR5 is 0.34 µg/ml. 2. Immobilized human FGF1 at 10 µg/ml (100 µl/well) can bind mouse FGFR5 with a linear range of 0.08-2 µg/ml. 3. Immobilized human bFGF at 5 µg/ml (100 µl/well) can bind mouse FGFR5. The EC50 of mouse FGFR5 is 0.22 µg/ml.

Properties

Purity	> 90 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 90 % as determined by reducing SDS-PAGE.

Background

Fibroblast growth factor receptor-like 1 (FGFR1) also known as Fibroblast growth factor receptor 5 (FGFR5), is a member of the fibroblast growth factor receptor (FGFR) family, where amino acid sequence is highly conserved between

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: www.elabscience.com

Tel: 1-832-243-6086

Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Recombinant Mouse FGFR1/FGFR5 Protein (His Tag)

Catalog Number:PKSM040824



members and throughout evolution. A full-length representative protein would consist of an extracellular region, composed of three immunoglobulin-like domains, a single hydrophobic membrane-spanning segment and a cytoplasmic tyrosine kinase domain. The extracellular portion of the protein interacts with fibroblast growth factors, setting in motion a cascade of downstream signals, ultimately influencing mitogenesis and differentiation. A unique feature of FGFR1/FGFR5 is that it does not contain an intracellular tyrosine kinase domain. Some muscle types, including the muscles of the tongue and the diaphragm, express FGFR1/FGFR5 at relatively high level. In contrast, the heart and the skeletal muscles of the limbs, as well as many other organs (brain, lung, liver, kidney, gut) express Fgfr1 only at basal level. It is conceivable that FGFR1/FGFR5 interacts with other Fgfrs, which are expressed in cartilage and muscle, to modulate FGF signaling.

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Tel: 1-832-243-6086

Fax: 1-832-243-6017

Web: www.elabscience.com

Email: techsupport@elabscience.com