EIF4A2 Polyclonal Antibody

Catalog Number: E-AB-19148



Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Reactivity Human, Mouse, Rat

Immunogen Fusion protein of human EIF4A2

Host Rabbit
Isotype IgG

Purification Antigen affinity purification


Conjugation Unconjugated

Formulation PBS with 0.05% NaN3 and 40% Glycerol,pH7.4


Applications Recommended Dilution

IHC 1:50-1:200 ELISA 1:5000-1:10000

Data



Immunohistochemistry of paraffin-embedded Human liver cancer tissue using EIF4A2 Polyclonal Antibody at dilution of 1:65(×200)



Immunohistochemistry of paraffin-embedded Human esophagus cancer tissue using EIF4A2 Polyclonal Antibody at dilution of 1:65(×200)

Preparation & Storage

Storage Store at -20°C. Avoid freeze / thaw cycles.

Background

ATP-dependent RNA helicase which is a subunit of the eIF4F complex involved in cap recognition and is required for mRNA binding to ribosome. In the current model of translation initiation, eIF4A unwinds RNA secondary structures in the 5'-UTR of mRNAs which is necessary to allow efficient binding of the small ribosomal subunit, and subsequent scanning for the initiator codon.

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: www.elabscience.com
Email: techsupport@elabscience.com