

Recombinant Human DMBT1/Muclin Protein (His Tag)

Catalog No. PKSH031031

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	GP340;muclin
Species	Human
Expression Host	HEK293 Cells
Sequence	Met 1-Ser 220
Accession	NP_004397.2
Calculated Molecular Weight	22.6 kDa
Observed molecular weight	35-45 kDa
Tag	C-His
Bioactivity	Immobilized recombinant human Galectin-3 at 10 µg/ml (100 µl/well) can bind biotinylated DMBT1-His with a linear range of 0.06-1.0 µg/ml.

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Deleted in malignant brain tumors 1 protein, also known as glycoprotein 340, surfactant pulmonary-associated D-binding

For Research Use Only

protein, DMBT1 and GP340, is a secreted protein which belongs to the DMBT1 family. DMBT1 contains 2 CUB domains, 14 SRCR domains and 1 ZP domain. It is highly expressed in alveolar and macrophage tissues. In some macrophages, expression is detected on the membrane, and in other macrophages, it is strongly expressed in the phagosome/phagolysosome compartments. Defects in DMBT1 are involved in the development of glioma (GLM). Gliomas are central nervous system neoplasms derived from glial cells and comprise astrocytomas, glioblastoma multiforme, oligodendrogliomas, and ependymomas. DMBT1 may be considered as a candidate tumor suppressor for brain, lung, esophageal, gastric, and colorectal cancers. It may play roles in mucosal defense system, cellular immune defense and epithelial differentiation. DMBT1 may play a role as an opsonin receptor for SFTPD and SPAR in macrophage tissues throughout the body, including epithelial cells lining the gastrointestinal tract. It may be an important factor in fate decision and differentiation of transit-amplifying ductular (oval) cells within the hepatic lineage. DMBT1 may function as a binding protein in saliva for the regulation of taste sensation. It binds to HIV-1 envelope protein and has been shown to both inhibit and facilitate viral transmission.