A Reliable Research Partner in Life Science and Medicine

## **Recombinant Mouse CCDC47 Protein (His Tag)**

Catalog No. PKSM040285

Note: Centrifuge before opening to ensure complete recovery of vial contents.

### **Description**

**Synonyms** 2610204L23Rik;asp4;C88307;calumin

**Species** Mouse

HEK293 Cells **Expression Host** Met1-Ser135 Sequence Accession NP\_080285.2 Calculated Molecular Weight 14.7 kDa C-His Tag

**Bioactivity** Not validated for activity

## **Properties**

**Purity** > 95 % as determined by reducing SDS-PAGE.

**Endotoxin** < 1.0 EU per µg of the protein as determined by the LAL method.

Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to **Storage** 

-80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots

of reconstituted samples are stable at < -20°C for 3 months.

**Shipping** This product is provided as lyophilized powder which is shipped with ice packs.

Formulation Lyophilized from sterile PBS, pH 7.4


Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as

protectants before lyophilization.

Please refer to the specific buffer information in the printed manual.

Reconstitution Please refer to the printed manual for detailed information.

#### Data



> 95 % as determined by reducing SDS-PAGE.

# **Background**

CCDC47 gene is expressed at high level. The gene contains 16 distinct gt-ag introns. Transcription produces 9 different mRNAs, 6 alternatively spliced variants and 3 unspliced forms. There are 3 probable alternative promotors, 3 non overlapping alternative last exons and 8 validated alternative polyadenylation sites. The mRNAs appear to differ by

#### For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Email: techsupport@elabscience.com

Web: www.elabscience.com

## **Elabscience Bionovation Inc.**



A Reliable Research Partner in Life Science and Medicine

truncation of the 5' end, truncation of the 3' end, presence or absence of a cassette exon, overlapping exons with different boundaries. Functionally, CCDC47 gene has been proposed to participate in processes such as calcium ion homeostasis, embryo development, ER overload response and post-embryonic development. CCDC47 are expected to have molecular function (calcium ion binding) and to localize in various compartments (membrane, endoplasmic reticulum, integral to membrane, microsome).

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Email: techsupport@elabscience.com

Web: www.elabscience.com