Caspase-8 Polyclonal Antibody

Catalog Number:E-AB-63533



Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Reactivity	Human,Mouse,Rat
Immunogen	Recombinant fusion protein of human Caspase-8 (NP_001073593.1).
Host	Rabbit
Isotype	IgG
Purification	Affinity purification
Conjugation	Unconjugated
Formulation	PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
Applications	Recommended Dilution
WB	1:500-1:1000
Data	



Western blot analysis of extracts of various cell lines using Caspase-8 Polyclonal Antibody at dilution of 1:1000. Observed Mw:55kDa

Calculated Mw:25-32kDa/45-61kDa

Preparation & Storage

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Background

This gene encodes a member of the cysteine-aspartic acid protease (caspase) family. Sequential activation of caspases plays a central role in the execution-phase of cell apoptosis. Caspases exist as inactive proenzymes composed of a prodomain, a large protease subunit, and a small protease subunit. Activation of caspases requires proteolytic processing at conserved internal aspartic residues to generate a heterodimeric enzyme consisting of the large and small subunits. This protein is involved in the programmed cell death induced by Fas and various apoptotic stimuli. The N-terminal FADD-like death effector domain of this protein suggests that it may interact with Fas-interacting protein FADD. This protein was detected in the insoluble fraction of the affected brain region from Huntington disease patients but not in those from normal controls, which implicated the role in neurodegenerative diseases. Many alternatively spliced transcript variants encoding different isoforms have been described, although not all variants have had their full-length sequences determined.

For Research Use Only

A Reliable Research Partner in Life Science and Medicine
Toll-free: 1-888-852-8623 Tel: 1-832-243-6086
Web: www.elabscience.com Email: techsupport@elabscience.com