

Recombinant Human TRAIL R1 / DR4 / TNFRSF10A (C-6His[®])



Catalog Number:PKSH034011

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	APO2;CD261 antigen;CD261;cytotoxic TRAIL receptor;DR4;TNF-related apoptosis-inducing ligand receptor 1;TNFRSF10A;TRAIL-R1;TRAILR1
Species	Human
Expression Host	HEK293 Cells
Sequence	Ala24-Asn239
Accession	O00220
Calculated Molecular Weight	23.7 kDa
Observed molecular weight	19-30 kDa
Tag	C-His

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Tumor necrosis factor receptor superfamily member 10A (TNFRSF10A) is also known as TNF-related apoptosis-inducing ligand receptor 1 (TRAIL-R1), Death receptor 4 (DR4), CD261 and APO2, which belongs to TNF superfamily. TNFRSF10A / DR4 is widely expressed and high levels are found in spleen, peripheral blood leukocytes, small intestine and thymus, but also in K-562 erythroleukemia cells, MCF-7 breast carcinoma cells and activated T-cells. APO2 / TNFRSF10A is receptor for the cytotoxic ligand TNFSF10 / TRAIL. This receptor is activated by tumor necrosis factor-related apoptosis inducing ligand (TNFSF1/TRAIL), and thus transduces cell death signal and induces cell apoptosis. TRAIL R1 can promote the activation of NF-kappa-B. TRAIL R1/CD261/TNFRSF1A induces apoptosis of many

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: www.elabscience.com

Tel: 1-832-243-6086

Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Recombinant Human TRAIL R1 / DR4 / TNFRSF10A (C-6His[®])



Catalog Number:PKSH034011

transformed cell lines but not of normal tissues, even though its death domain-containing receptor, DR4, is expressed on both cell types.

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: www.elabscience.com

Tel: 1-832-243-6086

Email: techsupport@elabscience.com

Fax: 1-832-243-6017