# **Recombinant Human PTPN11 protein (His tag)**

#### Catalog No. PDEH100261

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Synonyms	BPTP3;CFC;JMML;METCDS;NS1;PTN11;PTP-1D;PTP2C;SAP2;SH-PTP2;SH-PTP3;SHP2;SHPTP2;SHPTP3;Syp
Species	Human
Expression Host	E.coli
Sequence	Thr 2-Ala 521
Accession	Q06124
Calculated Molecular Weight	57.1 kDa
Observed molecular weight	60 kDa
Tag	N-His
Bioactivity	Not validated for activity
Properties	
Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	Please contact us for more information.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	It is recommended that sterile water be added to the vial to prepare a stock solution of 0.5 mg/mL. Concentration is measured by UV-Vis

### Data



> 95 % as determined by reducing SDS-PAGE.

## Background

#### **For Research Use Only**

# **Elabscience**®

SHP-2 (PTPN11) is a ubiquitously expressed, nonreceptor protein tyrosine phosphatase (PTP). It participates in signaling events downstream of receptors for growth factors, cytokines, hormones, antigens, and extracellular matrices in the control of cell growth, differentiation, migration, and death. Activation of SHP-2 and its association with Gab1 is critical for sustained Erk activation downstream of several growth factor receptors and cytokines. In addition to its role in Gab1-mediated Erk activation, SHP-2 attenuates EGF-dependent PI3 kinase activation by dephosphorylating Gab1 at p85 binding sites. SHP-2 becomes phosphorylated at Tyr542 and Tyr580 in its carboxy terminus in response to growth factor receptor activation. These phosphorylation events are thought to relieve basal inhibition and stimulate SHP-2 tyrosine phosphatase activity. Mutations in the corresponding gene result in a pair of clinically similar disorders (Noonan syndrome and LEOPARD syndrome) that may result from abnormal MAPK regulation.

For Research Use Only