B2M Polyclonal Antibody

Catalog Number:D-AB-10197L




Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description	
Reactivity	Human
Immunogen	Recombinant Human B2M protein expressed by E.coli
Host	Rabbit
Isotype	IgG
Purification	Antigen Affinity Purification
Conjugation	Unconjugated
Formulation	PBS with 0.02% sodium azide,50% glycerol pH 7.4
Applications	Recommended Dilution
WB	1:500-1:1000
IHC	1:100-1:200
Data	




Western blot with B2M Polyclonal antibody at dilution of 1:1000.lane 1:Hela whole cell lysate,lane 22A431 whole cell lysate,lane32Raji whole cell lysate Observed Mw:14kDa Calculated Mw:14kDa



Immunohistochemistry of paraffin-embedded Human colon using B2M Polycloanl Antibody at dilution of 1:100(100×)



Immunohistochemistry of paraffin-embedded Human colon using B2M Polycloanl Antibody at dilution of 1:100(400×)



Immunohistochemistry of paraffin-embedded Human tonsil using B2M Polycloanl Antibody at dilution of 1:200


For Research Use Only

A Reliable Research Partner in Life Science and Medicine Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Web: www.elabscience.com Email: techsupport@elabscience.com

B2M Polyclonal Antibody

Catalog Number:D-AB-10197L





Immunohistochemistry of paraffin-embedded Human kidney using B2M Polycloanl Antibody at dilution of 1:200

Preparation & Storage	
Storage	Store at -20°C. Avoid freeze / thaw cycles.
Background	l de la constante de

Beta-2-microglobulin (B2M) is a component of MHC class I molecules, which are present on the surface of nearly all nucleated cells. It can be found in body fluids under physiologic conditions as a result of shedding from cell surfaces or intracellular release. B2M has various biological functions, including antigen presentation. Investigations reveal that increased synthesis and release of B2M are present in several malignant diseases.

For Research Use Only