
## **IL2** Polyclonal Antibody

Catalog No. E-AB-40275


Note: Centrifuge before opening to ensure complete recovery of vial contents.

| Description  |  |
|--------------|--|
| Reactivity   | Human,Mouse  |
| Immunogen    | Recombinant Mouse Interleukin-2 protein                                  |
| Host         | Rabbit   |
| Isotype      | IgG  |
| Purification | Affinity Purification  |
| Conjugation  | Unconjugated   |
| Buffer       | PBS with 0.02% sodium azide,1% protective protein and 50% glycerol,pH7.4 |
| Applications | Recommended Dilution   |
| IHC          | 1:100-1:200  |
| Doto         |  |


Data



Immunohistochemistry of paraffin-embedded Human liver using IL2 Polyclonal Antibody at dilution of 1:100



Immunohistochemistry of paraffin-embedded Mouse liver using IL2 Polyclonal Antibody at dilution of 1:100



Immunohistochemistry of paraffin-embedded Human breast cancer using Il2 Polyclonal Antibody at dilution of 1:100

### **Preparation & Storage**

Storage

Store at -20°C. Avoid freeze / thaw cycles.

### Background

IL2 is a secreted cytokine involved in the operation and regulation of immune system. IL2 plays a pivotal role in T-cell

### **For Research Use Only**

Toll-free: 1-888-852-8623 Web: <u>www.elabscience.com</u>

# **Elabscience**®

activation and effector functions, including T cell proliferation, interferon-γproduction and cytotoxicity. IL2 can stimulate the propagation of lymphocytes and induce the cytotoxic T lymphocytes and lymphokine-activated killer cells for tumor cells. IL2 also influences memory T cell homeostasis through the regulation of memory T cell numbers and enhances major histocompatibility complex II (MHC II) expression on tumor cells.

For Research Use Only