

## Recombinant Human NLK Protein (His & GST Tag)

Catalog No. PKSH030360

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Synonyms</b>	LAK1;NLK
<b>Species</b>	Human
<b>Expression Host</b>	Baculovirus-Insect Cells
<b>Sequence</b>	Val121-Glu 527
<b>Accession</b>	Q9UBE8
<b>Calculated Molecular Weight</b>	74.1 kDa
<b>Observed molecular weight</b>	73 kDa
<b>Tag</b>	N-His-GST
<b>Bioactivity</b>	The specific activity was determined to be 3 nmol/min/mg using MBP as substrate.

### Properties

<b>Purity</b>	> 91 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Store at < -20°C, stable for 6 months. Please minimize freeze-thaw cycles.
<b>Shipping</b>	This product is provided as liquid. It is shipped at frozen temperature with blue ice/gel packs. Upon receipt, store it immediately at < -20°C.
<b>Formulation</b>	Supplied as sterile solution of 20mM Tris, 500mM NaCl, pH 8.0, 10% glycerol
<b>Reconstitution</b>	Not Applicable

### Data



> 91 % as determined by reducing SDS-PAGE.

### Background

Nemo-like kinase contains 1 protein kinase domain and belongs to the protein kinase superfamily, CMGC Ser/Thr protein kinase family and MAP kinase subfamily. It also contains a TQE activation loop motif in which autophosphorylation of the threonine residue (Thr-298) is sufficient for kinase activation. As a serine/threonine-protein kinase, nemo-like kinase regulates a number of transcription factors with key roles in cell fate determination. It is a positive effector of the non-canonical Wnt signaling pathway, acting downstream of WNT5A, MAP3K7/TAK1 and HIPK2. Activation of this pathway causes binding to and phosphorylation of the histone methyltransferase SETDB1. The NLK-SETDB1 complex

### For Research Use Only

subsequently interacts with PPARG, leading to methylation of PPARG target promoters at histone H3K9 and transcriptional silencing. The resulting loss of PPARG target gene transcription inhibits adipogenesis and promotes osteoblastogenesis in mesenchymal stem cells (MSCs). Nemo-like kinase also is a negative regulator of the canonical Wnt/beta-catenin signaling pathway.