Recombinant Human DOPA Decarboxylase/DDC Protein (His Tag)

Catalog No. PKSH030408

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description		
Synonyms	AADC	
Species	Human	
Expression Host	Baculovirus-Insect Cells	
Sequence	Met 1-Glu 480	
Accession	NP_000781.1	
Calculated Molecular Weight	55.0 kDa	
Observed molecular weight	48 kDa	
Tag	C-His	
Bioactivity	Measured by its ability to convert the substrate 3, 4-dihydroxy L-phenylalanine (L-Dopa) to 3, 4-dihydroxyphenylethylamine (dopamine). The dopamine product is measured by its absorbance at 340 nm after derivatization with trinitrobenzene sulfonic acid. The specific activity is > 1000 pmoles/min/ μ g.	
Properties		
Purity	> 90 % as determined by reducing SDS-PAGE.	
Endotoxin	< 1.0 EU per μ g of the protein as determined by the LAL method.	
Storage	Store at $< -20^{\circ}$ C, stable for 6 months. Please minimize freeze-thaw cycles.	
Shipping	This product is provided as liquid. It is shipped at frozen temperature with blue ice/gel packs. Upon receipt, store it immediately at $< -20^{\circ}$ C.	
Formulation	Supplied as sterile solution of 50mM Tris, 100mM NaCl, pH 8, 10% glycerol	
Reconstitution	Not Applicable	
Data		

KDa	ΜK	R
116	-	
66.2	-	-
45.0	-	-
35.0	-	
25.0	-	
18.4	-	
14.4	-	

> 90 % as determined by reducing SDS-PAGE.

Background

Dopa Decarboxylase (DDC), also known as AADC and Aromatic-L-amino acid decarboxylase, is a 54 kDa member of the group II decarboxylase family of proteins. It is a vitamin B6-dependent homodimeric enzyme that catalyzes the decarboxylation of both L-3,4-dihydroxyphenylalanine (L-DOPA) and L-5-hydroxytryptophan to dopamine and

For Research Use Only

Toll-free: 1-888-852-8623 Web: <u>www.elabscience.com</u> Tel: 1-832-243-6086 Email: <u>techsupport@elabscience.com</u>

Elabscience®

serotonin, respectively, which are major mammalian neurotransmitters and hormones belonging to catecholamines and indoleamines. Since L-DOPA is regularly used to treat the symptoms of Parkinson's disease, the catalytic pathway is of particular research interest. Defects of DDC are associated with severe developmental delay, oculogyric crises (OGC), as well as autosomal recessive disorder AADC deficiency, an early onset inborn error in neurotransmitter metabolism which can lead to catecholamine and serotonin deficiency.

For Research Use Only