Recombinant Human LILRB4/CD85k/ILT3 Protein (His Tag)

Catalog No. PKSH030439

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description		
Synonyms	CD85K;ILT-3;ILT3;LIR-5;LIR5	
Species	Human	
Expression Host	HEK293 Cells	
Sequence	Met 1-Glu259	
Accession	AAH26309.1	
Calculated Molecular Weight	27.7 kDa	
Tag	C-His	
Properties		
Purity	> 95 % as determined by reducing SDS-PAGE.	
Storage	Lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.	
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.	
Formulation	Lyophilized from sterile PBS, pH 7.4	
Reconstitution	Please refer to the printed manual for detailed information.	
Data		

KDa	MK	R
116	-	
66.2	-	
45.0		1000
35.0	-	-
25.0	-	
18.4	-	
^w 14.4	abseier	ice.com

Background

Mouse Leukocyte Immunoglobulin-like Receptor Subfamily B Member 4 (LILRB4/CD85k/ILT3) is an approximately transmembrane glycoprotein that negatively regulates immune cell activation. Mouse LILRB4 consists of a 215 amino acid (aa) extracellular domain with two Ig-like domains, a 22 aa transmembrane segment, and a 75 aa cytoplasmic domain with 3 immunoreceptor tyrosine-based inhibitory motifs (ITIM). Within the ECD, mouse LILRB4 shares 45% and 77% aa sequence identity with human and rat LILRB4, respectively. Alternative splicing of mouse LILRB4 generates a potentially soluble isoform that lacks the transmembrane segment. LILRB4 is expressed on dendritic cells (DC), monocytes, macrophages, and vascular endothelial cells (EC). Ligation of LILRB4 triggers ITIM-mediated inhibition of cellactivating signaling, leading to enhanced immune tolerance and reduced allogeneic graft rejection. Soluble LILRB4 induces the differentiation of CD8+ T suppressor cells (Ts) that can inhibit the effector functions of CD4+ Th cells and

For Research Use Only

Toll-free: 1-888-852-8623 Web: <u>www.elabscience.com</u>

Elabscience®

CD8+ CTL. In turn, CD8+ Ts cells induce LILRB4 up-regulation and a tolerogenic phenotype in monocytes, DC, and EC.

For Research Use Only