

Recombinant Human OBCAM/OPCML Protein (His Tag)

Catalog No. PKSH030473

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	IGLON1;OBCAM;OPCM
Species	Human
Expression Host	HEK293 Cells
Sequence	Met 1-Asn 322
Accession	NP_002536.1
Calculated Molecular Weight	34.0 kDa
Observed molecular weight	55-60 kDa
Tag	C-His
Bioactivity	Not validated for activity

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Opioid-binding Cell Adhesion Molecule (OBCAM), also known as OPCML, is a GPI-anchored cell adhesion molecule in the plasma membrane. This neuron-specific protein, consists of three immunoglobulin (Ig)-like domains anchored to the

For Research Use Only

membrane through a glycosylphosphatidylinositol (GPI)-tail. OPCML also belongs to the member of the IgLON family, a subgroup of the immunoglobulin superfamily, consisting of three members, LAMP, OBCAM, and Neurotrimin. These molecules interact homophilically and heterophilically within the family, and OBCAM acts only as heterodimers with LAMP or Neurotrimin and possibly inhibits neurite outgrowth from cerebellar granule cells. OBCAM has been presumed to play a role as a cell adhesion/recognition molecule. Furthermore, the OPCML protein defects may play an important role in the carcinogenesis of cervical or ovarian cancers, and this gene is regarded as a candidate TSG (tumor suppressor gene).