A Reliable Research Partner in Life Science and Medicine

# **Recombinant Human BPHL Protein (His Tag)**

Catalog No. PKSH030575

*Note:* Centrifuge before opening to ensure complete recovery of vial contents.

### **Description**

Synonyms BPH-RP;MCNAA;VACVASE

SpeciesHumanExpression HostE.coli

Sequence Ser38-Gln291
Accession Q86WA6-1
Calculated Molecular Weight 30.7 kDa
Observed molecular weight 31 kDa
Tag N-His

**Bioactivity** Not validated for activity

### **Properties**

**Purity** > 95 % as determined by reducing SDS-PAGE.

**Endotoxin** Please contact us for more information.

**Storage** Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to

-80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots

of reconstituted samples are stable at < -20°C for 3 months.

**Shipping** This product is provided as lyophilized powder which is shipped with ice packs.

Formulation Lyophilized from sterile 20mM Tris, 50mM NaCl, 10% glycerol, pH 8.0


Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as

protectants before lyophilization.

Please refer to the specific buffer information in the printed manual.

**Reconstitution** Please refer to the printed manual for detailed information.

#### Data



> 95 % as determined by reducing SDS-PAGE.

## **Background**

BPHL is a member of the serine protease family. BPHL is expressed large quantities in liver and kidney and in minor quantities in heart, intestine and skeletal muscle. BPHL is a specific alpha-amino acid ester hydrolase that prefers small,

#### For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: www.elabscience.com

Email: techsupport@elabscience.com

## **Elabscience Bionovation Inc.**



A Reliable Research Partner in Life Science and Medicine

hydrophobic, and aromatic side chains and does not have a stringent requirement for the leaving group other than preferring a primary alcohol. It catalyzes the hydrolytic activation of amino acid ester prodrugs of nucleoside analogs such as valacyclovir and valganciclovir. BPHL also activates valacyclovir to acyclovir. It may play a role in detoxification processes.

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Email: techsupport@elabscience.com

Web: www.elabscience.com