

# Recombinant Human MSN/Moesin Protein (aa 1-346, His Tag)



Catalog Number:PKSH030581

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


## Description

<b>Synonyms</b>	HEL70
<b>Species</b>	Human
<b>Expression Host</b>	E.coli
<b>Sequence</b>	Met 1-Glu 346
<b>Accession</b>	P26038
<b>Calculated Molecular Weight</b>	42.8 kDa
<b>Observed molecular weight</b>	45 kDa
<b>Tag</b>	N-His

## Properties

<b>Purity</b>	> 80 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	Please contact us for more information.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from sterile 20mM Tris, 0.5M NaCl, pH 8.0 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

## Data



> 80 % as determined by reducing SDS-PAGE.

## Background

Moesin is a member of the ERM family which includes ezrin and radixin. ERM proteins, highly related members of the larger protein 4.1 superfamily, can exist in an active or inactive conformation. It seems that ERM proteins function as cross-linkers between plasma membranes and actin-based cytoskeletons. The sole *Drosophila* ERM protein, moesin, functions to promote cortical actin assembly and apical-basal polarity. As a result, cells lacking moesin lose epithelial characteristics and adopt invasive migratory behaviour. It is localized to filopodia and other membranous protrusions that are important for cell-cell recognition and signaling and for cell movement. Moesin contains 1 FERM domain and is expressed in all tissues and cultured cells studied. Moesin has been shown to interact with CD43, Neutrophil cytosolic factor 1, VCAM-1, Neutrophil cytosolic factor 4, ICAM3 and EZR.

## For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086

Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017