

Recombinant Human AFM/Afamin Protein (His Tag)

Catalog No. PKSH030684

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	ALB2;ALBA;ALF
Species	Human
Expression Host	HEK293 Cells
Sequence	Met 1-Asn599
Accession	P43652
Calculated Molecular Weight	68.0 kDa
Observed molecular weight	68-94 kDa
Tag	C-His
Bioactivity	Not validated for activity

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Afamin is an 87 kDa glycoprotein with five predicted N-glycosylation sites. Afamin's glycan abundance contributes to conformational and chemical inhomogeneity presenting great challenges for molecular structure determination. Afamin, a

For Research Use Only

human plasma glycoprotein and putative transporter of hydrophobic molecules, has been shown to act as extracellular chaperone for poorly soluble, acylated Wnt proteins, forming a stable, soluble complex with functioning Wnt proteins. The 2.1-Å crystal structure of glycosylated human afamin reveals an almost exclusively hydrophobic binding cleft capable of harboring large hydrophobic moieties. Afamin plays a role in anti-apoptotic cellular processes related to oxidative stress and is associated with insulin resistance and other features of metabolic syndrome. Afamin may serve as a new early biomarker for pathological glucose metabolism during pregnancy. And first trimester screening for pre-eclampsia could be provided by a combination of afamin and placental bed vascularization. Moreover, the combination of first trimester serum afamin levels with BMI could provide a possible screening for gestational diabetes mellitus.