Recombinant Human TSC22D1 Protein (His Tag)

Catalog No. PKSH030789

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description		
Synonyms	Ptg-2;TGFB1I4;TSC22	
Species	Human	
Expression Host	E.coli	
Sequence	Met 1-Ala 144	
Accession	Q15714-2	
Calculated Molecular Weight	17.2 kDa	
Observed molecular weight	20 kDa	
Tag	N-His	
Bioactivity	Not validated for activity	
Properties		
Purity	> 92 % as determined by reducing SDS-PAGE.	
Endotoxin	Please contact us for more information.	
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.	
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.	
Formulation	Lyophilized from sterile PBS, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.	
Reconstitution	Please refer to the printed manual for detailed information.	

Data

KDa	MK	R
116	-	
66.2	-	
45.0	-	
35.0	-	
25.0	-	
18.4 14.4	=	

> 92 % as determined by reducing SDS-PAGE.

Background

TSC22 domain family, member 1 (TSC22D1) is one of the TGF-beta-stimulated clone-22 (TSC-22). TSC-22 was reported to be a differentiation-inducing factor which negatively regulates the growth of salivary gland cancer cells.

For Research Use Only

Toll-free: 1-888-852-8623 Web: <u>www.elabscience.com</u> Tel: 1-832-243-6086 Email: <u>techsupport@elabscience.com</u>

Elabscience®

TSC22D1, which encodes transforming growth factor beta-stimulated clone 22 (TSC-22), is thought to be a tumor suppressor because its expression is lost in many glioblastoma, salivary gland, and prostate cancers. TSC-22 is the founding member of the TSC-22/DIP/Bun family of leucine zipper transcription factors. TSC-22 may play an important role in maintaining the differentiated phenotype in salivary gland tumors, and may be a possible target of leukemia therapy. TSC22D1 forms homodimers via its conserved leucine zipper domain and heterodimerizes with TSC22D4. TSC22D1 has transcriptional repressor activity.

For Research Use Only