

Recombinant Human BVES/POPDC1 Protein (GST Tag)

Catalog No. PKSH030823

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	HBVES;POP1;POPDC1
Species	Human
Expression Host	E.coli
Sequence	Met 1-Asn36
Accession	NP_671488.1
Calculated Molecular Weight	30.8 kDa
Observed molecular weight	32.2 kDa
Tag	N-GST
Bioactivity	Not validated for activity

Properties

Purity	> 90 % as determined by reducing SDS-PAGE.
Endotoxin	Please contact us for more information.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 90 % as determined by reducing SDS-PAGE.

Background

Ferritin, light polypeptide (FTL) is the light subunit of the ferritin protein. Ferritin is the major intracellular iron storage protein in prokaryotes and eukaryotes. It is composed of 24 subunits of the heavy and light ferritin chains. Storage of iron

For Research Use Only

in the tissues occurs in the form of ferritin and hemosiderin. The latter originates from ferritin that has undergone intracellular digestion of its protein shell, leaving the iron core. Ferritin and hemosiderin are components of a continuum. Ferritin has been identified in all types of living organisms: animals, plants, molds, and bacteria. Within the protein shell of ferritin, iron is first oxidized to the ferric state for storage as ferric oxyhydroxide. Thus, ferritin removes excess iron from the cell sap where it could otherwise participate in peroxidation mechanisms.