

Recombinant Human SCGN/Secretagogin Protein (E.coli, His Tag)

Catalog No. PKSH030845

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	Secretagogin;SCGN;SECRET;CALBL;DJ501N12.8;setagin
Species	Human
Expression Host	E.coli
Sequence	Asp 2-Pro 276
Accession	NP_008929.2
Calculated Molecular Weight	33.4 kDa
Observed molecular weight	32 kDa
Tag	N-His
Bioactivity	Not validated for activity

Properties

Purity	> 97 % as determined by reducing SDS-PAGE.
Endotoxin	Please contact us for more information.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 97 % as determined by reducing SDS-PAGE.

Background

Secretagogin, also known as SCGN, is a secreted protein which is detectable in human serum after ischemic neuronal damage. It is a recently described calcium-binding protein. Secretagogin / SCGN is expressed at high levels in the

For Research Use Only

pancreatic islets of Langerhans and to a much lesser extent in the gastrointestinal tract (stomach, small intestine and colon), the adrenal medulla and cortex and the thyroid C-cells. In the brain, the expression of Secretagoin / SCGN is restricted to distinct subtypes of neurons with highest expression in the molecular layer of the cerebellum (stellate and basket cells), in the anterior part of the pituitary gland, in the thalamus, in the hypothalamus and in a subgroup of neocortical neurons. Secretagoin / SCGN is widely expressed in prostatic adenocarcinoma as opposed to adenocarcinomas in other organs. The function of Secretagoin / SCGN is unknown, but it has been suggested in beta-cells to influence calcium-influx and has been observed downregulated in diabetes-prone BB rat islets exposed to cytokines. Secretagoin / SCGN is involved in the calcium metabolism of tumour cells and endothelial cells in a subset of neoplasms of the brain and its coverings. Secretagoin / SCGN is also a novel marker for neuroendocrine differentiation.

For Research Use Only