Recombinant Human PROS1/Protein S Protein (His Tag)

Catalog No. PKSH030881

Note: Centrifuge before opening to ensure complete recovery of vial contents.

ynonyms	PROS;Protein S;PS21;PS22;PS23;PS24;PS25;PSA;THPH5;THPH6	
pecies	Human	
xpression Host	HEK293 Cells	
equence	Met 1-Ser676	
ccession	P07225	
alculated Molecular Weight	74.1 kDa	
bserved molecular weight	69-89 kDa	
ag	C-His	
ioactivity	Not validated for activity	
roperties		
urity	> 95 % as determined by reducing SDS-PAGE.	
ndotoxin	< 1.0 EU per μ g of the protein as determined by the LAL method.	
orage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.	
nipping	This product is provided as lyophilized powder which is shipped with ice packs.	
ormulation	Lyophilized from sterile PBS, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.	
econstitution	Please refer to the printed manual for detailed information.	

Data

KDa	MK	R
116	-	-
66.2	-	
45.0	-	
35.0	-	
25.0	-	
18.4 14.4	=	
. 4.4		

> 95 % as determined by reducing SDS-PAGE.

Background

PROS1, also known as protein S, is a vitamin K-dependent plasma protein that functions as a cofactor for the anticoagulant protease, activated protein C (APC) to inhibit blood coagulation. PROS1 has two isoforms: a free,

For Research Use Only

Toll-free: 1-888-852-8623 Web: <u>www.elabscience.com</u> Tel: 1-832-243-6086 Email: <u>techsupport@elabscience.com</u>

Elabscience®

functionally active form and an inactive form complexed with C4b-binding protein. Besides its anticoagulant function, PROS1 also acts as an agonist for the tyrosine kinase receptors Tyro3, Axl, and Mer. The endothelium expresses Tyro3, Axl, and Mer and produces protein S. The interaction of protein S with endothelial cells and particularly its effects on angiogenesis have not yet been analyzed.