Recombinant Human TWF1/Twinfilin-1 Protein (His & GST Tag)

Catalog No. PKSH030992

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Synonyms	A6;MGC23788;MGC41876;PTK9
Species	Human
Expression Host	E.coli
Sequence	Met 1-Asp 252
Accession	Q12792-4
Calculated Molecular Weight	57.0 kDa
Observed molecular weight	50 kDa
Tag	N-His-GST
Bioactivity	Not validated for activity

Properties

Purity	> 84 % as determined by reducing SDS-PAGE.
Endotoxin	Please contact us for more information.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80℃. Reconstituted protein solution can be stored at 4-8℃ for 2-7 days. Aliquots of reconstituted samples are stable at < -20℃ for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile 50mM Tris, 500mM NaCl, pH 8.0 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 84 % as determined by reducing SDS-PAGE.

Background

Twinfilin-1; also known as Protein A6; Protein tyrosine kinase 9; TWF1 and PTK9; is a cytoplasm protein which belongs to theactin-binding proteins ADF family and Twinfilin subfamily. Twinfilin-1 (TWF1 / PTK9) is a highly conserved

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017
Web: www.elabscience.com Email: techsupport@elabscience.com

actin monomer-binding protein that regulates cytoskeletal dynamics in organisms from yeast to mammals. In addition to the mammalian twinfilin-1; a second protein with approximately 65% sequence identity to twinfilin-1 exists in mouse and humans. TWF1 / PTK9 is expressed at high levels in the colon; testis; ovary; prostate and lung. It is expressed at lower levels in the brain; bladder and heart. It is not detected in liver. TWF1 / PTK9 is an actin-binding protein involved in motile and morphological processes. It inhibits actin polymerization; likely by sequestering G-actin. By capping the barbed ends of filaments; it also regulates motility. TWF1 / PTK9 seems to play an important role in clathrin-mediated endocytosis and distribution of endocytic organelles.

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017
Web: www.elabscience.com Email: techsupport@elabscience.com