

# Recombinant Human SIRP gamma/CD172g Protein (His Tag)



Catalog Number:PKSH030993

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


## Description

<b>Synonyms</b>	Signal-Regulatory Protein Gamma;SIRP-Gamma;CD172 Antigen-Like Family Member B;Signal-Fegulatory Protein Beta-2;SIRP-b2;SIRP-Beta-2;CD172g;SIRPG;SIRPB2
<b>Species</b>	Human
<b>Expression Host</b>	HEK293 Cells
<b>Sequence</b>	Met 1-Ser 364
<b>Accession</b>	NP_061026.2
<b>Calculated Molecular Weight</b>	34.7 kDa
<b>Tag</b>	C-His
<b>Bioactivity</b>	Measured by its binding ability in a functional ELISA. Immobilized human SIRPG-His at 10 µg/ml (100 µl/well) can bind human CD47-Fc, The EC50 of human CD47-Fc is 0.58-1.34 µg/ml.

## Properties

<b>Purity</b>	> 96 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from sterile PBS, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

## Data



> 96 % as determined by reducing SDS-PAGE.

## Background

Signal-regulatory protein gamma (SIRPG/SIRP gamma) also known as CD172 antigen-like family member B, CD172g, and CD172g antigen, is a member of the signal-regulatory protein (SIRP) family, and also belongs to the immunoglobulin superfamily. SIRP family members are receptor-type transmembrane glycoproteins known to be involved in the negative regulation of receptor tyrosine kinase-coupled signaling processes. SIRPG/SIRP gamma/CD172g is probable

## For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086

Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017

# Recombinant Human SIRP gamma/CD172g Protein (His Tag)



Catalog Number:PKSH030993

---

immunoglobulin-like cell surface receptor. On binding with CD47, SIRPG can mediate cell-cell adhesion. SIRPG/SIRP gamma is engagement on T-cells by CD47 on antigen-presenting cells results in enhanced antigen-specific T-cell proliferation and costimulates T-cell activation. SIRPG/SIRP gamma/CD172g is detected in liver, and at very low levels in brain, heart, lung, pancreas, kidney, placenta and skeletal muscle. Expressed on CD4+ T-cells, CD8+ T-cells, CD56-bright natural killer (NK) cells, CD20+ cells, and all activated NK cells. This cytokine is mainly present in the paracortical T-cell area of lymph nodes, with only sparse positive cells in the mantle and in the germinal center of B-cell follicles. In the thymus, SIRPG is primarily expressed in the medulla on mature T-lymphocytes that have undergone thymic selection.

---

## For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Tel: 1-832-243-6086

Fax: 1-832-243-6017

Web: [www.elabscience.com](http://www.elabscience.com)

Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)