

Recombinant Human USP7/HAUSP Protein (aa 208-560)

Catalog Number:PKSH031028



Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	HAUSP;TEF1
Species	Human
Expression Host	Baculovirus-Insect Cells
Sequence	Lys 208-Glu 560
Accession	NP_003461.2
Calculated Molecular Weight	41.0 kDa
Observed molecular weight	41 kDa
Tag	None

Properties

Purity	> 98 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile 50mM Tris, 100mM NaCl, 0.5mM PMSF, pH 8.0 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 98 % as determined by reducing SDS-PAGE.

Background

Ubiquitin carboxyl-terminal hydrolase 7, also known as Ubiquitin thioesterase 7, Herpesvirus-associated ubiquitin-specific protease, Ubiquitin-specific-processing protease 7, USP7 and HAUSP, is a widely expressed protein which belongs to the peptidase C19 family. USP7 is a member of the family of deubiquitinating enzymes. It is involved in the regulation of stress response pathways, epigenetic silencing and the progress of infections by DNA viruses. USP7 is a protein with a cysteine peptidase core, N- and C-terminal domains required for protein-protein interactions. USP7 contributes to epigenetic silencing of homeotic genes by Polycomb (Pc). USP7 cleaves ubiquitin fusion protein substrates. It deubiquitinates TP53/p53 and MDM2 and strongly stabilizes TP53 even in the presence of excess MDM2. USP7 also induces TP53-dependent cell growth repression and apoptosis. USP7 has key roles in the p53 pathway whereby it

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: www.elabscience.com

Tel: 1-832-243-6086

Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Recombinant Human USP7/HAUSP Protein (aa 208-560)

Catalog Number:PKSH031028



stabilizes both p53 and MDM2. Herpes simplex virus type 1 (HSV-1) regulatory protein ICP0 stimulates lytic infection and the reactivation of quiescent viral genomes. ICP0 interacts very strongly with USP7. USP7-mediated stabilization of ICP0 is dominant over ICP0-induced degradation of USP7 during productive HSV-1 infection. The biological significance of the ICP0-USP7 interaction may be most pronounced in natural infection situations, in which limited amounts of ICP0 are expressed.

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: www.elabscience.com

Tel: 1-832-243-6086

Email: techsupport@elabscience.com

Fax: 1-832-243-6017