A Reliable Research Partner in Life Science and Medicine

## **Recombinant Human B3GNT2 Protein (Fc Tag)**

Catalog No. PKSH031125

Note: Centrifuge before opening to ensure complete recovery of vial contents.

### **Description**

Synonyms B3GN-T2;B3GNT;B3GNT-2;B3GNT;BETA3GNT;BGnT-2;BGNT2

Species Human

Expression Host HEK293 Cells
Sequence Lys29-Cys397
Accession Q9NY97-1
Calculated Molecular Weight 71.2 kDa
Observed molecular weight 112-120 kDa
Tag N-hFc

**Bioactivity** Not validated for activity

### **Properties**

**Purity** > 90 % as determined by reducing SDS-PAGE.

Endotoxin < 1.0 EU per ug of the protein as determined by the LAL method.

**Storage** Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to

-80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots

of reconstituted samples are stable at < -20°C for 3 months.

**Shipping** This product is provided as lyophilized powder which is shipped with ice packs.

**Formulation** Lyophilized from sterile PBS, pH 7.4


Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as

protectants before lyophilization.

Please refer to the specific buffer information in the printed manual.

**Reconstitution** Please refer to the printed manual for detailed information.

#### Data



> 90 % as determined by reducing SDS-PAGE.

### **Background**

B3GNT2 belongs to the beta-1,3-N-acetylglucosaminyltransferase family. It is a type II transmembrane protein which prefers the substrate of lacto-N-neotetraose. Alternative splicing produced 2 isoforms of the human protein. B3GNT2

#### For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: www.elabscience.com

Email: techsupport@elabscience.com

# **Elabscience Bionovation Inc.**



A Reliable Research Partner in Life Science and Medicine

catalyzes the initiation and elongation of poly-N- acetyllactosamine chains. Enzymatic activities of some glycosyltransferases are markedly increased via complex formation with other transferases or cofactor proteins. B3GNT2 and beta3Gn-T8 can form a heterodimer in vitro and that the complex exhibits much higher enzymatic activity than either enzyme alone. It is found that up-regulation of beta3Gn-T8 in differentiated HL-60 cells may increases poly-Nacetyllactosamine chains by activating intrinsic B3GNT2.

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Email: techsupport@elabscience.com

Web: www.elabscience.com