A Reliable Research Partner in Life Science and Medicine

Recombinant Human RHEB Protein (His Tag)

Catalog No. PKSH031137

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Synonyms GTP-Binding Protein Rheb;Ras Homolog Enriched in Brain;RHEB;RHEB2

Species Human

Expression Host Baculovirus-Insect Cells

SequenceMet 1-Cys181AccessionQ15382Calculated Molecular Weight22.4 kDaObserved molecular weight21 kDaTagN-His

Bioactivity Not validated for activity

Properties

Purity > 85 % as determined by reducing SDS-PAGE.

Endotoxin < 1.0 EU per ug of the protein as determined by the LAL method.

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to

-80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots

of reconstituted samples are stable at < -20°C for 3 months.

Shipping This product is provided as lyophilized powder which is shipped with ice packs.

Formulation Lyophilized from sterile 20mM Tris, 500mM NaCl, 10% glycerol, pH 7.4


Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as

protectants before lyophilization.

Please refer to the specific buffer information in the printed manual.

Reconstitution Please refer to the printed manual for detailed information.

Data



> 85 % as determined by reducing SDS-PAGE.

Background

RHEB is a recently discovered member of the Ras superfamily that may be involved in neural plasticity. This function is novel and not typically associated with the Ras proteins. RHEB gene is a member of the small GTPase superfamily and

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: www.elabscience.com

Email: techsupport@elabscience.com

Elabscience Bionovation Inc.



A Reliable Research Partner in Life Science and Medicine

encodes a lipid-anchored; cell membrane protein with five repeats of the RAS-related GTP-binding region. RHEB is vital in regulation of growth and cell cycle progression due to its role in the insulin / TOR / S6K signaling pathway. The protein has GTPase activity and shuttles between a GDP-bound form and a GTP-bound form; and farnesylation of RHEB is required for this activity. Three pseudogenes have been mapped; two on chromosome 10 and one on chromosome 22.

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Email: techsupport@elabscience.com

Web: www.elabscience.com