Recombinant Human EphA4 Protein (His & Fc Tag)

Catalog No. PKSH031153

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Synonyms	Ephrin type-A receptor 4;HEK8;SEK;TYRO1;EPHA4;Tyrosine-protein kinase receptor SEK;Tyrosine-protein kinase TYRO1;EK8;hEK8;EPH-like kinase 8
Species	Human
Expression Host	HEK293 Cells
Sequence	Met 1-Thr 547
Accession	NP_004429.1
Calculated Molecular Weight	86.5 kDa
Observed molecular weight	100-110 kDa
Tag	C-His-Fc
Bioactivity	Immobilized human EPHA5 at 20 μ g/ml (100 μ l/well) can bind human EFNA4-Fc with a linear ranger of 1. 28-32 ng/ml.
Properties	
Properties Purity	> 92 % as determined by reducing SDS-PAGE.
Properties Purity Endotoxin	 > 92 % as determined by reducing SDS-PAGE. < 1.0 EU per μg of the protein as determined by the LAL method.
Properties Purity Endotoxin Storage	 > 92 % as determined by reducing SDS-PAGE. < 1.0 EU per μg of the protein as determined by the LAL method. Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Properties Purity Endotoxin Storage Shipping	 > 92 % as determined by reducing SDS-PAGE. < 1.0 EU per µg of the protein as determined by the LAL method. Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months. This product is provided as lyophilized powder which is shipped with ice packs.
Properties Purity Endotoxin Storage Shipping Formulation	 > 92 % as determined by reducing SDS-PAGE. < 1.0 EU per µg of the protein as determined by the LAL method. Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months. This product is provided as lyophilized powder which is shipped with ice packs. Lyophilized from sterile PBS, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Properties Purity Endotoxin Storage Shipping Formulation Reconstitution	 > 92 % as determined by reducing SDS-PAGE. < 1.0 EU per µg of the protein as determined by the LAL method. Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months. This product is provided as lyophilized powder which is shipped with ice packs. Lyophilized from sterile PBS, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.

Data



> 92 % as determined by reducing SDS-PAGE.

Background

For Research Use Only

Elabscience®

EPH receptor A4 (ephrin type-A receptor 4), also known as EphA4, belongs to the ephrin receptor subfamily of the protein-tyrosine kinase family which 16 known receptors (14 found in mammals) are involved: EPHA1, EPHA2, EPHA3, EPHA4, EPHA5, EPHA6, EPHA7, EPHA8, EPHA9, EPHA10, EPHB1, EPHB2, EPHB3, EPHB4, EPHB5, EPHB6. The Eph family of receptor tyrosine kinases (comprising EphA and EphB receptors) has been implicated in synapse formation and the regulation of synaptic function and plasticity6. EphA4 is enriched on dendritic spines of pyramidal neurons in the adult mouse hippocampus, and ephrin-A3 is localized on astrocytic processes that envelop spines. Eph receptor–mediated signaling, which is triggered by ephrins7, probably modifies the properties of synapses during synaptic activation and remodeling. Ephrin receptors are components of cell signalling pathways involved in animal growth and development, forming the largest sub-family of receptor tyrosine kinases (RTKs). The extracellular domain of an EphA4 interacts with ephrin ligands, which may be tethered to neighbouring cells. Ligand-mediated activation of Ephs induce various important downstream effects and Eph receptors have been studied for their potential roles in the development of cancer.

For Research Use Only