Recombinant Human TPST1 Protein (His Tag)

Catalog No. PKSH031177

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description		
Synonyms	TANGO13A	
Species	Human	
Expression Host	HEK293 Cells	
Sequence	Gln 26-Glu 370	
Accession	NP_003587.1	
Calculated Molecular Weight	41.7 kDa	
Observed molecular weight	45-48 kDa	
Tag	N-His	
Bioactivity	Not validated for activity	
Properties		
Purity	> 80 % as determined by reducing SDS-PAGE.	
Endotoxin	< 1.0 EU per μ g of the protein as determined by the LAL method.	
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.	
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.	
Formulation	Lyophilized from sterile PBS, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.	
Reconstitution	Please refer to the printed manual for detailed information.	

Data

KDa	МК	R
116	Entre	-
66.2	-	dence
45.0	-	-
35.0	-	-1er
		Elabson
25.0	-	
18.4	e clente	
14.4	-	

> 80 % as determined by reducing SDS-PAGE.

Background

Protein-tyrosine sulfotransferase 1, also known as Tyrosylprotein sulfotransferase 1 and TPST1, is a single-pass type II membrane proteinwhich belongs to the protein sulfotransferase family. Tyrosine O-sulfation is a common posttranslational

For Research Use Only

Toll-free: 1-888-852-8623 Web: <u>www.elabscience.com</u> Tel: 1-832-243-6086 Email: <u>techsupport@elabscience.com</u>

Elabscience®

modification of proteins in all multicellular organisms. This reaction is mediated by a Golgi enzyme activity called tyrosylprotein sulfotransferase (TPST) that catalyzes the transfer of sulfate from 3'-phosphoadenosine 5'-phosphosulfate to tyrosine residues within acidic motifs of polypeptides. Tyrosine O-sulfation has been shown to be important in protein-protein interactions in several systems. Tyrosine sulfation is mediated by one of two Golgi isoenzymes, called tyrosylprotein sulfotransferases (TPST-1 and TPST-2). A relatively small number of proteins are known to undergo tyrosine sulfation, including certain adhesion molecules, G-protein-coupled receptors, coagulation factors, serpins, extracellular matrix proteins, and hormones. TPST1 is a human tyrosylprotein sulfotransferase that uses 3'phosphoadenosine-5'phosphosulfate (PAPS) to transfer the sulfate moiety to proteins predominantly designated for secretion. TPST1 bears N-linked glycosyl residues exclusively at position Asn60 and Asn262. TPST1 and TPST2 have distinct biological roles that may reflect differences in their macromolecular substrate specificity.