

## Recombinant Human B4GALT1/GGTB2 Protein (His Tag)

**Catalog No.** PKSH031198

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


### Description

<b>Synonyms</b>	B4GAL-T1;beta4Gal-T1;CDG2D;GGTB2;GT1;GTB
<b>Species</b>	Human
<b>Expression Host</b>	HEK293 Cells
<b>Sequence</b>	Gly 44-Ser 398
<b>Accession</b>	NP_001488.2
<b>Calculated Molecular Weight</b>	41.5 kDa
<b>Observed molecular weight</b>	45-55 kDa
<b>Tag</b>	N-His
<b>Bioactivity</b>	Not validated for activity

### Properties

<b>Purity</b>	> 92 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU per µg of the protein as determined by the LAL method.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from sterile PBS, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

### Data



> 92 % as determined by reducing SDS-PAGE.

### Background

Beta1,4-Galactosyltransferase-I (B4GALT1), one of seven beta1,4-galactosyltransferases, is an enzyme commonly found in the trans-Golgi complex that adds galactose to oligosaccharides. They have an N-terminal hydrophobic signal sequence

### For Research Use Only

that directs the protein to the Golgi apparatus and which then remains uncleaved to function as a transmembrane anchor. By sequence similarity, the beta4GalTs form four groups: beta4GalT1 and beta4GalT2, beta4GalT3 and beta4GalT4, beta4GalT5 and beta4GalT6, and beta4GalT7. B4GALT1 gene directs production of B4GALT1 protein using either of two transcription start sites. The product of the smaller transcript serves the traditional biosynthetic role in the Golgi. This form also complexes with  $\alpha$ -lactalbumin, a mammary-specific protein, to form lactose synthase. In addition to a biosynthetic role, the protein translated from the longer transcript appears on the plasma membranes of some cells where it serves as a signalling receptor in cell-matrix interactions such as sperm-egg binding.