

# Recombinant Human PRDM2/RIZ1 Protein (GST Tag)

Catalog Number:PKSH031204



**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


## Description

<b>Synonyms</b>	HUMHOXY1;KMT8;MTB-ZF;RIZ;RIZ1;RIZ2
<b>Species</b>	Human
<b>Expression Host</b>	E.coli
<b>Sequence</b>	Met 1-Ala 200
<b>Accession</b>	NP_036363.2
<b>Calculated Molecular Weight</b>	49.6 kDa
<b>Observed molecular weight</b>	49.6 kDa
<b>Tag</b>	N-GST

## Properties

<b>Purity</b>	> 86 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	Please contact us for more information.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from sterile 20mM Tris, 150mM NaCl, 0.5mM DTT, 0.5mM GSH, pH 8.0 Normally 5 % - 8 % trehalose, mannitol and 0.01 % Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed ma
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

## Data



> 86 % as determined by reducing SDS-PAGE.

## Background

PR domain containing 2; with ZNF domain (PRDM2); also known as zinc finger protein RIZ; is a member of histone methyltransferase (HMT) class enzymes that methylate lysine residues of histones or proteins. HMTs contain a conserved catalytic core termed the SET domain; which shares sequence homology with an independently described sequence motif; the PR domain. PRDM2 contains 8 C2H2-type zinc fingers and a distinct SET domain; and is highly expressed in retinoblastoma cell lines and in brain tumors; as well as in a number of other cell lines and in brain; heart; skeletal muscle; liver and spleen. PRDM2 is a S-adenosyl-L-methionine-dependent histone methyltransferase that specifically methylates 'Lys-9' of histone H3; and is identified as a tumor suppressor. It is reported that intact PR( SET) sequence is required for

## For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086

Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017

# Recombinant Human PRDM2/RIZ1 Protein (GST Tag)

Catalog Number:PKSH031204



tumor suppression functions; mutations in the PR domain caused activity reduction in human cancers. Also; S-adenosylhomocysteine or methyl donor deficiency inhibits RIZ1 and other H3 lysine 9 methylation activities. PRDM2 may also function as a DNA-binding transcription factor. It Binds to the macrophage-specific TPA-responsive element (MTE) of the HMOX1 (heme oxygenase 1) gene and act as a transcriptional activator. In addition; PRDM2 (RIZ) is able to binds to the retinoblastoma protein (RB) and also Interacts with GATA3.

---

## For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Tel: 1-832-243-6086

Fax: 1-832-243-6017

Web: [www.elabscience.com](http://www.elabscience.com)

Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)