

Recombinant Human ENTPD5 Protein (His Tag)

Catalog No. PKSH031224

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	CD39L4;NTPDase-5;PCPH
Species	Human
Expression Host	Baculovirus-Insect Cells
Sequence	Met 1-His 428
Accession	O75356
Calculated Molecular Weight	47.0 kDa
Observed molecular weight	45 kDa
Tag	C-His
Bioactivity	Measured by its ability to hydrolyze the 5'phosphate groups from the substrate guanosine5'diphosphate (GDP). The specific activity is > 7, 000 pmoles/min/μg.

Properties

Purity	> 90 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per μg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile 20mM Tris, 500mM NaCl, pH 7.4, 10% glycerol Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 90 % as determined by reducing SDS-PAGE.

Background

Ectonucleoside triphosphate diphosphohydrolase 5 (ENTPD5), also known as CD39 antigen-like 4, ER-UDPase,

For Research Use Only

Guanosine-diphosphatase ENTPD5, Nucleoside diphosphatase Uridine-diphosphatase ENTPD5. This hydrolase is expressed in response to phosphoinositide 3-kinase (PI3K) signaling. Activation of PI3K results in FOXO phosphorylation by AKT1 and loss of ENTPD5 transcriptional repression. It is Up-regulated in PTEN-deficient cells. Uridine diphosphatase (UDPase) that promotes protein N-glycosylation and ATP level regulation. ENTPD5 promotes protein N-glycosylation and folding in the endoplasmic reticulum, as well as elevated ATP consumption in the cytosol via an ATP hydrolysis cycle. Together with CMPK1 and AK1, ENTPD5 constitutes an ATP hydrolysis cycle that converts ATP to AMP and results in a compensatory increase in aerobic glycolysis. ENTPD5 also hydrolyzes GDP and IDP but not any other nucleoside di-, mono- or triphosphates, nor thiamine pyrophosphate. This enzyme Plays a key role in the AKT1-PTEN signaling pathway by promoting glycolysis in proliferating cells in response to phosphoinositide 3-kinase (PI3K) signaling.

For Research Use Only