

PE Anti-Human CD192/CCR2 Antibody[K036C2]

Catalog No.	E-AB-F1385D	Reactivity	Human
Storage	Store at 2~8°C, Avoid freeze / thaw cycles	Applications	FCM

Important Note: Centrifuge before opening to ensure complete recovery of vial contents.

Antigen Information

Alternate Names	CKR2, CCR2A, CCR2B, CKR2A, CKR2B, CMKBR2, MCP-1-R, CC-CKR-2
Uniprot ID	P41597
Gene ID	729230
Background	CCR2 is a chemokine receptor that binds monocyte chemoattractant proteins (MCP-1, 2, 3 and 4). Two spliced variants were initially described for CCR2 (CCR2A and CCR2B). These variants differ in their terminal carboxyl tails. Monocyte adhesion to the arterial endothelium and subsequent migration into the intima are central events in the pathogenesis of atherosclerosis. CCR2 and MCP-1 have been associated to atherosclerotic plaques. MCP-1 is induced by modified-LDL in endothelial cells and may trigger firm adhesion of monocytes to vascular endothelium under flow conditions. Local overexpression of MCP-1 at vessel walls induces infiltration of macrophages and formation of atherosclerotic lesions. Obesity induces an inflammatory state that is implicated in many clinically important complications, including insulin resistance, diabetes, atherosclerosis, and non-alcoholic fatty liver disease. CCR2 influences the development of obesity and associated adipose tissue inflammation.

Product Details

Form	Liquid
Size	20Tests/100Tests/100Tests×2
Clone No.	K036C2
Host	Mouse
Isotype	Mouse IgG2a, κ
Reactivity	Human
Application	FCM
Isotype Control	PE Mouse IgG2a, κ Isotype Control[C1.18.4] [Product E-AB-F09802D]
Storage Buffer	Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.
Shipping	Biological ice pack at 4 °C
Stability & Storage	Keep as concentrated solution. Store at 2~8°C and protected from prolonged exposure to light. Do not freeze. This product is guaranteed up to one year from purchase.

For Research Use Only

Fluorophore


Conjugation: PE

PE is designed to be excited by the Blue (488 nm), Green (532 nm) and Yellow-Green (561 nm) lasers and detected using an optical filter centered near 575 nm (e.g., a 585/42 nm bandpass filter).

Recommended usage

Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. The amount of the reagent is suggested to be used 5 µL of antibody per test (million cells in 100 µL staining volume or per 100 µL of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

Product data



Staining of normal human peripheral blood cells with PE Anti-Human CD192/CCR2 Antibody[K036C2] (filled gray histogram) or PE Mouse IgG2a, κ Isotype Control (empty black histogram). Cells in the monocytes gate were used for analysis.

Related Information

1. Sample Preparation for Flow Cytometry <https://www.elabscience.com/List-detail-5594.html>
2. Staining Cell Surface Targets for Flow Cytometry <https://www.elabscience.com/List-detail-5568.html>
3. Flow Cytometry Troubleshooting Tips <https://www.elabscience.com/List-detail-5593.html>
4. How to select the appropriate detection channel through the spectrogram? <https://www.elabscience.com/List-detail-459742.html>