

Recombinant Human NAALADL1 Protein (His Tag)

Catalog No. PKSH031237

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	I100;NAALADASEL;NAALADL1
Species	Human
Expression Host	HEK293 Cells
Sequence	Pro 29-Leu 740
Accession	NP_005459.2
Calculated Molecular Weight	80.0 kDa
Observed molecular weight	90 kDa
Tag	N-His
Bioactivity	Not validated for activity

Properties

Purity	> 97 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 97 % as determined by reducing SDS-PAGE.

Background

N-acetylated-alpha-linked acidic dipeptidase-like protein, also known as NAALADL1, NAALADase L, and Ileal dipeptidylpeptidase, is a Single-pass type I I membrane protein and a member of the peptidase M28 family and M28B

For Research Use Only

subfamily. NAALADase L is mainly expressed in the distal small intestine. It is also expressed in the spleen and testis and Weakly expressed in the brain, locating mainly to the brain stem, amygdala, thalamus and ventral striatum. NAALADase L is a chloride-activated, membrane bound, metallopeptidase that cleaves the endogenous neuropeptide N-acetyl-aspartyl-glutamate (NAAG). NAAG acts as a partial NMDA agonist as well as a full agonist at the presynaptic metabotropic glutamate receptor 3 (mGluR3), where it acts to reduce glutamate release. NAALADase L also exhibits a dipeptidyl-peptidase IV type activity. NAALADase inhibition may be a novel therapeutic approach to reduce or inhibit heightened aggressiveness, and possibly to treat aggressive behavior associated with psychiatric disorders.