Recombinant Human S100A10 Protein (His Tag)

Catalog No. PKSH031240

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Synonyms	42C;ANX2L;ANX2LG;CAL1L;Ca[1];CLP11;GP11;MGC111133;p10;P11
Species	Human
Expression Host	E.coli
Sequence	Pro 2-Lys 97
Accession	NP_002957.1
Calculated Molecular Weight	12.6 kDa
Observed molecular weight	12.6 kDa
Tag	N-His
Bioactivity	Not validated for activity
Properties	
Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	Please contact us for more information.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile 20mM Tris, pH 8.5, 10% glycerol Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

S100 proteinis a family of low molecular weight protein found in vertebrates characterized by twoEF-hand calciumbinding motifs. There are at least 21 different S100 proteins, and the name is derived from the fact that the protein

For Research Use Only

Toll-free: 1-888-852-8623 Web: <u>www.elabscience.com</u> Tel: 1-832-243-6086 Email: <u>techsupport@elabscience.com</u>

Elabscience®

is100% soluble in ammonium sulfateat neutralpH. S100 proteins have been implicated in a variety of intracellular and extracellular functions. They are involved in regulation of protein phosphorylation, transcription factors, the dynamics of cytoskeleton constituents, enzyme activities, cell growth and differentiation, and the inflammatory response. Protein S100-A10, also known as Calpactin I light chain, Cellular ligand of annexin II, ANX2LG and S100A10, is a member of the S100 family. In contrast to all other S100 proteins, S100A10 is Ca(2+) insensitive because of amino acid replacements in its Ca(2+)-binding loops that lock the protein in a permanently active state. S100A10 forms a heterotetramer with annexin IIH and promotes carcinoma invasion and metastasis by plasminogen activation. S100A10 and annexin II contribute to the aggressive characteristics of anaplastic carcinoma, while playing a constitutive role in papillary carcinoma. S100A10 induces the dimerization of ANXA2 / p36, it may function as a regulator of protein phosphorylation in that the ANXA2 monomer is the preferred target of tyrosine-specific kinase. S100A10 functions as a linker tethering certain transmembrane proteins to annexin A2 thereby assisting their traffic to the plasma membrane and/or their firm anchorage at certain membrane sites.

For Research Use Only