## Recombinant Human CCD5L Protein (His Tag)

### Catalog No. PKSH031429

*Note:* Centrifuge before opening to ensure complete recovery of vial contents.

Description		
Synonyms	AIM;API6;PRO229;SP-ALPHA;Spalpha	
Species	Human	
Expression Host	HEK293 Cells	
Sequence	Met 1-Gly 347	
Accession	NP_005885.1	
Calculated Molecular Weight	37.5 kDa	
Observed molecular weight	45 kDa	
Tag	C-His	
Bioactivity	Not validated for activity	
Properties		
Purity	> 97 % as determined by reducing SDS-PAGE.	
Endotoxin	< 1.0 EU per $\mu$ g of the protein as determined by the LAL method.	
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.	
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.	
Formulation	Lyophilized from sterile PBS, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.	
Reconstitution	Please refer to the printed manual for detailed information.	

Data

KDa	MK	R
116 66.2	-	
45.0 35.0	-	-
25.0	-	
18.4 14.4	=	

> 97 % as determined by reducing SDS-PAGE.

## Background

CD5L, also known as CD5 antigen-like, is a soluble protein belonging to group B of the scavenger receptor cysteine-rich (SRCR) superfamily and contains three SRCR domains. It is a secreted glycoprotein and expressed by macrophages

#### **For Research Use Only**

Toll-free: 1-888-852-8623 Web: <u>www.elabscience.com</u> Tel: 1-832-243-6086 Email: <u>techsupport@elabscience.com</u>

# **Elabscience**®

presentin lymphoid tissues (spleen, lymph node, thymus, and bone marrow). It binds to myelomonocytic and lymphoid cells and may play an important role in the regulation of the innate and adaptive immune systems. CD5L functions as a pattern recognition molecule by binding both lipoteichoic acid (LTA) on Gram positive and lipopolysaccharide (LPS) on Gram negative bacteria. and the SRCR domain 1 of CD5L retains both the LPS and LTA binding activities. In addition, it is revealed that CD5L seems to play a role as an inhibitor of apoptosis.

For Research Use Only