Recombinant Human CD30/TNFRSF8 Protein (His Tag)

Catalog Number: PKSH031431



Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description		
Synonyms	Tumor necrosis factor receptor superfamily member 8;CD30L receptor;Ki-1 antigen;Lymphocyte activation antigen CD30;CD30;TNFRSF8;D1S166E	
Species	Human	
Expression Host	HEK293 Cells	
Sequence	Met 1-Lys 379	
Accession	NP_001234.2	
Calculated Molecular Weight	40 kDa	
Observed molecular weight	75-90 kDa	
Tag	C-His	
Bioactivity	Measured by its binding ability in a functional ELISA. 2. Immobilized recombinant human CD30 at 2 μ g/ml (100 μ l/well) can bind human CD30L Fc Chimera with a linear range of 6. 4-800 ng/ml.	
Properties		
Purity	> 95 % as determined by reducing SDS-PAGE.	
Endotoxin	< 1.0 EU per μ g of the protein as determined by the LAL method.	
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.	
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.	
Formulation	Lyophilized from sterile PBS, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.	
Reconstitution	Please refer to the printed manual for detailed information.	
Data		

KDa	MK	R
116	-	-
66.2	-	10.
45.0	-	
35.0	-	
25.0	-	
18.4	-	
14.4	-	

> 95 % as determined by reducing SDS-PAGE.

Background

CD30, also known as TNFRSF8, is a cell membrane protein of the tumor necrosis factor receptor (TNFR) superfamily. CD30 protein is expressed by activated, but not resting, T and B cells. CD30 can regulate proliferation of lymphocytes and may also play an important role in human immunodeficiency virus replication. As a regulator of apoptosis, CD30 protein induces cell death or proliferation, depending on the cell type, and has been shown to limit the proliferative

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Recombinant Human CD30/TNFRSF8 Protein (His Tag)

Catalog Number:PKSH031431



potential of autoreactive CD8 effector T cells and protect the body against autoimmunity. CD30 protein expression is upregulated in various hematological malignancies, including Reed-Sternberg cells in Hodgkin's disease (HD), anaplastic large cell lymphoma (ALCL) and subsets of Non-Hodgkin's lymphomas (NHLs), and CD30 is also linked to leukocytes in patients with chronic inflammatory diseases, including lupus erythematosus, asthma, rheumatoid arthritis and atopic dermatitis (AD).

For Research Use Only

A Reliable Research Partner in Life Science and Medicine Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Web: www.elabscience.com Email: techsupport@elabscience.com