

Recombinant Human MICB Protein (His & Fc Tag)

Catalog No. PKSH031444

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	PERB11.2
Species	Human
Expression Host	HEK293 Cells
Sequence	Met 1-Gly 298
Accession	NP_005922.2
Calculated Molecular Weight	59.5 kDa
Observed molecular weight	80-90 kDa
Tag	C-His-Fc
Bioactivity	Immobilized human His-NKG2D (78-216) at 10 µg/ml (100 µl/well) can bind human MICB-Fc, The EC50 of human MICB-Fc is 15.9-37.1 ng/ml.

Properties

Purity	> 98 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 98 % as determined by reducing SDS-PAGE.

Background

MHC class I polypeptide-related sequence B; also known as MICB; is a heavily glycosylated protein serving as a ligand for

For Research Use Only

the type II receptor NKG2D. MICB shares 85% amino acid identity with MICA; a closely related protein; both of which contain three extracellular immunoglobulin-like domains; but without capacity to bind peptide or interact with beta-2-microglobulin. acting as a stress-induced self-antigen; binding of MICB to the NKG2D receptor activates the cytolytic response of natural killer (NK) cells; CD8+ $\alpha\beta$ T cells; and $\gamma\delta$ T cells on which the receptor is expressed. MICA/B are minimally expressed on normal cells; but are frequently expressed on epithelial tumors and can be induced by bacterial and viral infections. MICA/B recognition thus is involved in tumor surveillance; viral infections; and autoimmune diseases.