Recombinant Human MICB Protein (His & Fc Tag)

Catalog Number: PKSH031444



Note: Centrifuge before opening to ensure complete recovery of vial contents.

		- 40	
Desc	\mathbf{r}	$\mathbf{n}\mathbf{T}\mathbf{I}$	\mathbf{on}
DUSU	ш		ULL

Synonyms PERB11.2 Species Human

Expression Host
Sequence
Met 1-Gly 298
Accession
NP_005922.2
Calculated Molecular Weight
Observed molecular weight
Tag
HEK293 Cells
Met 1-Gly 298
NP_005922.2
Sequence
NP_005922.2
Calculated Molecular Weight
C-His-Fc

Bioactivity Immobilized human His-NKG2D (78-216) at 10 μg/ml (100 μl/well) can bind

human MICB-Fch, The EC50 of human MICB-Fch is 15.9-37.1 ng/ml.

Properties

Purity > 98 % as determined by reducing SDS-PAGE.

Endotoxin < 1.0 EU per µg of the protein as determined by the LAL method.

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to

-80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots

of reconstituted samples are stable at < -20°C for 3 months.

Shipping This product is provided as lyophilized powder which is shipped with ice packs.

Formulation Lyophilized from sterile PBS, pH 7.4


Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as

protectants before lyophilization.

Please refer to the specific buffer information in the printed manual.

Reconstitution Please refer to the printed manual for detailed information.

Data



> 98 % as determined by reducing SDS-PAGE.

Background

MHC class I polypeptide-related sequence B; also known as MICB; is a heavily glycosylated protein serving as a ligand for the type II receptor NKG2D. MICB shares 85% amino acid identity with MICA; a closely related protein; both of which contain three extracellular immunoglobulin-like domains; but without capacity to bind peptide or interact with beta-2-microglobulin. acting as a stress-induced self-antigen; binding of MICB to the NKG2D receptor activates the cytolytic response of natural killer (NK) cells; CD8+ $\alpha\beta$ T cells; and $\gamma\delta$ T cells on which the receptor is expressed. MICA/B are minimally expressed on normal cells; but are frequently expressed on epithelial tumors and can be induced

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: www.elabscience.com Email: techsupport@elabscience.com

Recombinant Human MICB Protein (His & Fc Tag)

Catalog Number:PKSH031444



by bacterial and viral infections. MICA/B recognition thus is involved in tumor surveillance; viral infections; and autoimmune diseases.

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: <u>www.elabscience.com</u> Email: <u>techsupport@elabscience.com</u>