# Recombinant Human PDK/PDZ binding kinase/TOPK Protein (His Tag)

by Elabscience

Catalog Number: PKSH031457

Note: Centrifuge before opening to ensure complete recovery of vial contents.

#### **Description**

Synonyms CT84;HEL164;Nori-3;SPK;TOPK

Species Human

**Expression Host** Baculovirus-Insect Cells

SequenceMet 1-Val 322AccessionNP\_060962.2Calculated Molecular Weight37.0 kDaObserved molecular weight37 kDaTagC-His

#### **Properties**

**Purity** > 80 % as determined by reducing SDS-PAGE.

**Endotoxin**  $< 1.0 \text{ EU per } \mu \text{g of the protein as determined by the LAL method.}$ 

**Storage** Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to

-80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots

of reconstituted samples are stable at < -20°C for 3 months.

**Shipping** This product is provided as lyophilized powder which is shipped with ice packs.

**Formulation** Lyophilized from sterile PBS, pH 7.4


Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as

protectants before lyophilization.

Please refer to the specific buffer information in the printed manual.

**Reconstitution** Please refer to the printed manual for detailed information.

#### Data



> 80 % as determined by reducing SDS-PAGE.

### **Background**

PDZ binding kinase (PBK), also known as TOPK (T-LAK cell-originated protein kinase), is a serine/threonine kinase related to the dual specific mitogen-activated protein kinase kinase (MAPKK) family, and has all the characteristic protein kinase subdomains and a C-terminal PDZ-binding T/SXV motif. PBK is expressed in the testis restrictedly expressed in outer cell layer of seminiferous tubules, as well as placenta. PBK may be enrolled in the activation of lymphoid cells and support testicular functions, with a suggested role in the process of spermatogenesis. This mitotic kinase phosphorylates MAP kinase p38 and seems to be active in mitosis. When phosphorylated, PBK forms a protein-protein interaction with tumor suppressor p53 (TP53), leading to TP53 destabilization and attenuation of G2/M checkpoint during doxorubicin-induced DNA damage. The expression level of PBK is thus upregulated in a variety of

#### For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: <u>www.elabscience.com</u> Email: <u>techsupport@elabscience.com</u>

# Recombinant Human PDK/PDZ binding kinase/TOPK **Protein (His Tag)**



Catalog Number:PKSH031457

neoplasms including hematological malignancies.

## For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017 Email: techsupport@elabscience.com

Web: www.elabscience.com