Recombinant Human PDGFRa/CD140a Protein (Fc Tag)

Catalog Number: PKSH031529



Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Synonyms CD140A;PDGFR-2;PDGFR2;RHEPDGFRA;Platelet-derived growth factor

> receptor alpha;PDGFR-alpha;Alpha platelet-derived growth factor receptor;CD140 antigen-like family member A;Platelet-derived growth factor alpha receptor;Platelet-

derived growth factor receptor 2

Species Human

Expression Host HEK293 Cells Met 1-Glu 524 **Sequence** Accession NP 006197.1 Calculated Molecular Weight 82.9 kDa 120-130 kDa Observed molecular weight C-hFc

Tag

Bioactivity 1. Measured by its ability to bind human PDGF-C in a functional ELISA.

2. Measured by its ability to bind mouse PDGF-C in a functional ELISA.

Properties

Purity > 90 % as determined by reducing SDS-PAGE.

Endotoxin < 1.0 EU per ug of the protein as determined by the LAL method.

Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to Storage

-80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots

of reconstituted samples are stable at < -20°C for 3 months.

Shipping This product is provided as lyophilized powder which is shipped with ice packs.

Formulation Lyophilized from sterile PBS, pH 7.4


Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as

protectants before lyophilization.

Please refer to the specific buffer information in the printed manual.

Reconstitution Please refer to the printed manual for detailed information.

Data



> 90 % as determined by reducing SDS-PAGE.

Background

PDGFRA; also known as CD140a; together with the structurally homolog protein PDGFRB (CD140b); are cell surface receptors for members of the platelet-derived growth factor family. They are members of the class III subfamily of receptor tyrosine kinase (RTKs) with the similar structure characteristics of five immunoglobulin-like domains in their

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: www.elabscience.com Email: techsupport@elabscience.com

Recombinant Human PDGFRa/CD140a Protein (Fc Tag)

Catalog Number: PKSH031529



extracellular region and a split kinase domain in their intracellular region. PDGFRA is expressed in oligodendrocyte progenitor cells and mesothelial cell; and binds all three ligand isoforms PDGF-AA; PDGF-BB and PDGF-AB with high affinity; whereas PDGFRB dose not bind PDGF-AA. PDGFRA plays an essential role in regulating proliferation; chemotaxis and migration of mesangial cells. Recent studies have indicated that PDGFRA acts as a critical mediator of signaling in testis organogenesis and Leydig cell differentiation; and in addition; particularly important for kidney development. Additionally; PDGFRA is involved in tumor angiogenesis and maintenance of the tumor microenvironment and has been implicated in development and metastasis of Hepatocellular carcinoma (HCC). PDGFRA may represent a potential therapeutic target in thymic tumours. PDGFRA gene amplification rather than gene mutation may be the underlying genetic mechanism driving PDGFRA overexpression in a portion of gliomas.

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: www.elabscience.com Email: techsupport@elabscience.com