Recombinant Human RELT/TNFRSF19L Protein (His Tag)

Catalog Number: PKSH031543



Note: Centrifuge before opening to ensure complete recovery of vial contents.

T .	.
LIACOPT	ntion
Descri	77717

Synonyms Tumor necrosis factor receptor superfamily member 19L;TNFRSF19L;Receptor

expressed in lymphoid tissues;RELT

Species Human

Expression Host HEK293 Cells
Sequence Met 1-Ala 160
Accession NP_116260.2
Calculated Molecular Weight 15.7 kDa
Tag C-His

Properties

Purity > 95 % as determined by reducing SDS-PAGE.

Endotoxin < 1.0 EU per μg of the protein as determined by the LAL method.

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to

-80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots

of reconstituted samples are stable at < -20°C for 3 months.

Shipping This product is provided as lyophilized powder which is shipped with ice packs.

Formulation Lyophilized from sterile PBS, pH 7.4


Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as

protectants before lyophilization.

Please refer to the specific buffer information in the printed manual.

Reconstitution Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Receptor expressed in lymphoid tissues (RELT); also known as tumor necrosis factor receptor superfamily; member 19-like (TNFRSF19L); is a member of the TNF-receptor superfamily. This receptor is especially abundant in hematologic tissues. It has been shown to activate the NF-kappaB pathway and selectively bind TNF receptor-associated factor 1. RELT/TNFRSF19L is capable of stimulating T-cell proliferation in the presence of CD3 signaling; which suggests its regulatory role in immune response. RELT/TNFRSF19L is a type I transmembrane glycoprotein with a cysteine-rich extracellular domain; possessing significant homology to other members of the TNFR superfamily; especially TNFRSF19; DR3; OX40; and LTbeta receptor. RELT/TNFRSF19L is able to activate the NF-kappaB pathway and selectively binds tumor necrosis factor receptor-associated factor 1. RELT/TNFRSF19L is able to activate the NF-κB pathway and selectively binds tumor necrosis factor receptor-associated factor 1. Although the soluble form of RELT fusion protein

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: www.elabscience.com Email: techsupport@elabscience.com

Recombinant Human RELT/TNFRSF19L Protein (His Tag)

Catalog Number:PKSH031543



does not inhibit the one-way mixed lymphocyte reaction; immobilized RELT/TNFRSF19L is capable of costimulating T-cell proliferation in the presence of CD3 signaling.

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: <u>www.elabscience.com</u> Email: <u>techsupport@elabscience.com</u>