

Recombinant Human TREM1 Protein (His & Fc Tag)

Catalog Number:PKSH031561



Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	Triggering Receptor Expressed on Myeloid Cells 1;TREM-1;Triggering Receptor Expressed on Monocytes 1;CD354;TREM1
Species	Human
Expression Host	HEK293 Cells
Sequence	Met 1-Arg 200
Accession	NP_061113.1
Calculated Molecular Weight	48.3 kDa
Observed molecular weight	60-65 kDa
Tag	C-His-Fc

Properties

Purity	> 97 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01 % Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 97 % as determined by reducing SDS-PAGE.

Background

TREM1 (triggering receptor expressed on myeloid cells) is a type I transmembrane protein with a single Ig-like domain; and is selectively expressed on blood neutrophils and a subset of monocytes. As a member of the growing family of receptors related to NK cell receptors; TREM1 activates downstream signaling events with the help of an adapter protein called DAP12. Expression of TREM1 is up-regulated by bacterial LPS; a ligand for TLR4; as well as lipoteichoic acid. Although its natural ligand has not been identified; engagement of TREM1 with agonist mAbs triggers secretion of the proinflammatory cytokines TNF- α and IL-1 β ; as well as chemokines such as IL-8 and monocyte chemoattractant protein (MCP)-1. Intracellularly; TREM1 induces Ca²⁺ mobilization and tyrosine phosphorylation of extracellular signal-related

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: www.elabscience.com

Tel: 1-832-243-6086

Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Recombinant Human TREM1 Protein (His & Fc Tag)

Catalog Number:PKSH031561



kinase 1 (ERK1); ERK2 and phospholipase C- γ . In an animal model of LPS-induced septic shock; blockade of TREM1 signaling inhibited hyperresponsiveness and death. Thus; it has been demonstrated that TREM1 performs a critical function in immune responses involved in host defense against microbial challenges; and is suggested to be a potential therapeutic target for septic shock.

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Tel: 1-832-243-6086

Fax: 1-832-243-6017

Web: www.elabscience.com

Email: techsupport@elabscience.com