

Recombinant Human IFNGR1 Protein (His Tag)

Catalog No. PKSH031670

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	CD119;IFNGR;IMD27A;IMD27B;Interferon gamma receptor 1;IFNGR1
Species	Human
Expression Host	HEK293 Cells
Sequence	Met 1-Gly 245
Accession	NP_000407.1
Calculated Molecular Weight	27.3 kDa
Tag	C-His
Bioactivity	Measured by its ability to inhibit rhIFN γ mediated protection of WISH cells infected with vesicular stomatitis virus(VSV). The ED50 for this effect is typically 2-8 μ g/mL.

Properties

Purity	> 97 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per μ g of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 97 % as determined by reducing SDS-PAGE.

Background

The cluster of differentiation (CD) system is commonly used as cell markers in immunophenotyping. Different kinds of

For Research Use Only

cells in the immune system can be identified through the surface CD molecules which associating with the immune function of the cell. There are more than 320 CD unique clusters and subclusters have been identified. Some of the CD molecules serve as receptors or ligands important to the cell through initiating a signal cascade which then alter the behavior of the cell. Some CD proteins do not take part in cell signal process but have other functions such as cell adhesion. CD119 (cluster of differentiation 119), also known as IFNGR1 (interferon gamma receptor 1), is part of the heterodimeric gamma interferon receptor which consists of IFNGR1 (CD119) and IFNGR2. The IFNGR1 gene encodes the ligand-binding chain (alpha) of the interteron receptor while IFNGR gene encodes the non-ligand binding partner. The ability of the interferon- γ was achieved through binding to the interferon receptor CD119. After binding, the products of activated T-lymphocytes interferon- γ exerts antiviral activity, growth inhibitory effect, and several immune- regulatory activities on a variety of cell types.