

Recombinant Human MERTK/MER Protein (His & Fc Tag)

Catalog Number:PKSH031700

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	Tyrosine-protein kinase Mer;Proto-oncogene c-Mer;Receptor tyrosine kinase MerTK;MERTK;MER;Mer
Species	Human
Expression Host	HEK293 Cells
Sequence	Met 1-Ala 499
Accession	NP_006334.2
Calculated Molecular Weight	80.0 kDa
Observed molecular weight	140-150 kDa
Tag	C-His-Fc

Properties

Purity	> 97 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01 % Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 97 % as determined by reducing SDS-PAGE.

Background

Proto-oncogene tyrosine-protein kinase MER (MERTK) is a member of the MER/AXL/TYRO3 receptor kinase family and encodes a transmembrane protein with two fibronectin type-III domains; two Ig-like C2-type (immunoglobulin-like) domains; and one tyrosine kinase domain. MERTK is localized in membrane and is not expressed in normal B- and T-lymphocytes but is expressed in numerous neoplastic B- and T-cell lines. This protein is highly expressed in testis; ovary; prostate; lung; and kidney; with lower expression in spleen; small intestine; colon; and liver. MERTK regulates many physiological processes including cell survival; migration; differentiation; and phagocytosis of apoptotic cells (efferocytosis). Ligand binding at the cell surface induces autophosphorylation of MERTK on its intracellular domain that

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: www.elabscience.com

Tel: 1-832-243-6086

Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Recombinant Human MERTK/MER Protein (His & Fc Tag)



Catalog Number:PKSH031700

provides docking sites for downstream signaling molecules. MERTK signaling plays a role in various processes such as macrophage clearance of apoptotic cells; platelet aggregation; cytoskeleton reorganization and engulfment. MERTK plays also an important role in inhibition of Toll-like receptors (TLRs)-mediated innate immune response by activating STAT1; which selectively induces production of suppressors of cytokine signaling SOCS1 and SOCS3. Defects in MERTK are the cause of retinitis pigmentosa type 38.

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: www.elabscience.com

Tel: 1-832-243-6086

Email: techsupport@elabscience.com

Fax: 1-832-243-6017