

Recombinant Human Noggin/NOG Protein (His Tag)

Catalog Number:PKSH031720



Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	Noggin;SYM1;SYNS1
Species	Human
Expression Host	HEK293 Cells
Sequence	Met 1-Cys 232
Accession	NP_005441.1
Calculated Molecular Weight	24.6 kDa
Observed molecular weight	30 kDa
Tag	C-His
Bioactivity	Measured by its ability to inhibit recombinant human BMP4-induced alkaline phosphatase production by MC3T3-E1 cells. The ED50 for this effect is typically 0.05-0.3 µg/mL in the presence of 50 ng/mL of recombinant human BMP4.

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01 % Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Noggin is a secreted protein involved at multiple stages of vertebrate embryonic development including neural induction and is known to exert its effects by inhibiting the bone morphogenetic protein (BMP)-signaling pathway. It binds several BMPs with very high (picomolar) affinities; with a marked preference for BMP2 and BMP4 over BMP7. By binding tightly to BMPs; Noggin prevents BMPs from binding their receptors. Noggin binds the bone morphogenetic proteins (BMP) such as BMP-4 and BMP-7; and inhibits BMP signaling by blocking the molecular interfaces of the binding

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: www.elabscience.com

Tel: 1-832-243-6086

Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Recombinant Human Noggin/NOG Protein (His Tag)

Catalog Number:PKSH031720



epitopes for both type I and type II receptors. Interaction of BMP and its antagonist Noggin governs various developmental and cellular processes; including embryonic dorsal-ventral axis; induction of neural tissue; formation of joints in the skeletal system and neurogenesis in the adult brain. Noggin plays a key role in neural induction by inhibiting BMP4; along with other TGF- β signaling inhibitors such as chordin and follistatin. Mouse knockout experiments have demonstrated that noggin also plays a crucial role in bone development; joint formation; and neural tube fusion.

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Tel: 1-832-243-6086

Fax: 1-832-243-6017

Web: www.elabscience.com

Email: techsupport@elabscience.com