
## Recombinant Human CD31/PECAM1 Protein (His Tag)

Catalog Number: PKSH031833



*Note:* Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Synonyms	Platelet endothelial cell adhesion molecule;PECAM-1;EndoCAM;GPIIA;PECA1;CD31;PECAM1
Species	Human
Expression Host	HEK293 Cells
Sequence	Met 1-Lys 601
Accession	EAW94208.1
Calculated Molecular Weight	66.0 kDa
Observed molecular weight	100-110 kDa
Tag	C-His
Bioactivity	Measured by the ability of the immobilized protein to support the adhesion of Jurkat human acute T cell leukemia cells. When $8 \times 104$ cells/well are added to CD31-His coated plates (5µg/mL, 100 µL/well) in the presence of 10 µg/mL PHA, approximately > 70% will adhere after 30 minutes at 37°C.
Properties	
Purity	> 97 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per $\mu$ g of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.
Data	



> 97 % as determined by reducing SDS-PAGE.

## Background

The Cluster of Differentiation 31 (CD31) adhesion molecule, also known as platelet-endothelial cell adhesion molecule-1 (PECAM-1), is the only known member of the CAM family on platelets. CD31 protein is a 130-kDa transmembrane glycoprotein expressed by endothelial cells, platelets, monocytes, neutrophils, and certain T cell subsets. CD31 protein is

## For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Catalog Number: PKSH031833



also expressed in certain tumors, including epithelioid hemangioendothelioma, other vascular tumors, and histiocytic malignancies. CD31 plays a key role in removing aged neutrophils and tissue regeneration. CD31 protein mediates the homotypic or heterotypic cell adhesion by binding to itself or the leukocyte integrin  $\alpha\nu\beta3$ , and thus plays a role in neutrophil recruitment in inflammatory responses, transendothelial migration of leukocytes, as well as in cardiovascular development.

## For Research Use Only

A Reliable Research Partner in Life Science and Medicine Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Web: www.elabscience.com Email: techsupport@elabscience.com