

Recombinant Human VEGF-A/VEGF121 Protein (His Tag)

Catalog No. PKSH031983

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	Vascular endothelial growth factor A;VEGF-A;Vascular permeability factor;VPF;VEGFA;VEGF
Species	Human
Expression Host	HEK293 Cells
Sequence	Ala27-Arg147
Accession	P15692-9
Calculated Molecular Weight	15.1 kDa
Observed molecular weight	16-18 kDa
Tag	C-His
Bioactivity	Not validated for activity

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg of the protein as determined by the LAL method.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Human VEGF121; also known as Vascular endothelial growth factor A; VEGFA; Vascular permeability factor; VPF and

For Research Use Only

VEGF; is a homodimeric; heparin-binding glycoprotein which belongs to the platelet-derived growth factor (PDGF)/vascular endothelial growth factor (VEGF) family. VEGF-A is a glycosylated mitogen that specifically acts on endothelial cells and has various effects; including mediating increased vascular permeability; inducing angiogenesis; vasculogenesis; permeabilization of blood vessels and endothelial cell growth; increasing microvascular permeability; promoting cell migration and inhibiting apoptosis. Alternatively spliced transcript variants of VEGF-A encode either secreted or cell-associated isoforms. The lymphangiogenesis may be promoted by upregulation of VEGF121; which may in turn act in part via induction of VEGF-C. It binds to the FLT1/VEGFR1 and KDR/VEGFR2 receptors; heparan sulfate and heparin. NRP1/Neuropilin-1 binds isoforms VEGF-165 and VEGF-145. Isoform VEGF165B binds to KDR but does not activate downstream signaling pathways; does not activate angiogenesis and inhibits tumor growth.